## Supplementary Information of

## Non-peptide compounds from Kromopolites svenhedini (Verhoeff) and their anti-tumor and iNOS inhibitory activities

Yuan-Nan Yuan<sup>1,2</sup>, Jin-Qiang Li<sup>2,3</sup>, Hong-Bin Fang<sup>2</sup>, Shao-Jun Xing<sup>3</sup>, Yong-Ming Yan<sup>2\*</sup>, and Yong-Xian Cheng<sup>1,2\*\*</sup>

Address: <sup>1</sup> School of Pharmacy, Guangdong Pharmaceutical University, Guangzhou 510006, PR China. <sup>2</sup> Institute for Inheritance-Based Innovation of Chinese Medicine, School of Pharmaceutical Sciences, Health Science Center, Shenzhen University, Shenzhen 518060, PR China. <sup>3</sup> Department of Pathogen Biology, Health Science Center, Shenzhen University, Shenzhen 518060, PR China.

Email: Yong-Ming Yan - yanym@szu.edu.cn; Yong-Xian Cheng yxcheng@szu.edu.cn

\* Corresponding author

## Contents

Figure S1. <sup>1</sup>H NMR spectrum of **1** in methanol-d<sub>4</sub>

- Figure S2. <sup>13</sup>C NMR and DEPT spectrum of **1** in methanol-*d*<sub>4</sub>
- Figure S3. <sup>1</sup>H–<sup>1</sup>H COSY spectrum of **1** in methanol-*d*<sub>4</sub>
- Figure S4. HSQC spectrum of 1 in methanol-d4
- Figure S5. HMBC spectrum of 1 in methanol-d4
- Figure S6. HRESIMS spectrum of 1
- Figure S7. CD spectrum of 1 in MeOH
- Figure S8. <sup>1</sup>H NMR spectrum of **2** in methanol-d<sub>4</sub>
- Figure S9. <sup>13</sup>C NMR and DEPT spectrum of 2 in methanol-d<sub>4</sub>
- Figure S10. <sup>1</sup>H–<sup>1</sup>H COSY spectrum of 2 in methanol-d<sub>4</sub>
- Figure S10. HSQC spectrum of **2** in methanol-*d*<sub>4</sub>
- Figure S12. HMBC spectrum of 2 in methanol-d4
- Figure S13. HRESIMS spectrum of 2
- Figure S14. <sup>1</sup>H NMR spectrum of **3** in methanol-d<sub>4</sub>
- Figure S15. <sup>13</sup>C NMR and DEPT spectrum of 3 in methanol-d<sub>4</sub>
- Figure S16. <sup>1</sup>H–<sup>1</sup>H COSY spectrum of **3** in methanol-*d*<sub>4</sub>
- Figure S17. HSQC spectrum of 3 in methanol-d4
- Figure S18. HMBC spectrum of 3 in methanol-d<sub>4</sub>
- Figure S19. HRESIMS spectrum of 3
- Figure S20. <sup>1</sup>H NMR spectrum of **4** in methanol-d<sub>4</sub>
- Figure S21. <sup>13</sup>C NMR and DEPT spectrum of **4** in methanol-d<sub>4</sub>
- Figure S22. <sup>1</sup>H–<sup>1</sup>H COSY spectrum of **4** in methanol-*d*<sub>4</sub>
- Figure S23. HSQC spectrum of 4 in methanol-d4
- Figure S24. HMBC spectrum of 4 in methanol-d4
- Figure S25. HRESIMS spectrum of 4
- Figure S26. CD spectrum of 4 in MeOH
- Figure S27. <sup>1</sup>H NMR spectrum of 7 in methanol-d<sub>4</sub>
- Figure S28. <sup>13</sup>C NMR and DEPT spectrum of 7 in methanol-d<sub>4</sub>
- Figure S29. <sup>1</sup>H–<sup>1</sup>H COSY spectrum of **7** in methanol-*d*<sub>4</sub>
- Figure S30. HSQC spectrum of 7 in methanol-d4
- Figure S31. HMBC spectrum of 7 in methanol-d<sub>4</sub>
- Figure S32. ROESY spectrum of 7 in methanol-d4
- Figure S33. HRESIMS spectrum of 7
- Figure S34. <sup>1</sup>H NMR spectrum of **8** in methanol-*d*<sub>4</sub>
- Figure S35. <sup>13</sup>C NMR and DEPT spectrum of 8 in methanol-d<sub>4</sub>
- Figure S36. <sup>1</sup>H–<sup>1</sup>H COSY spectrum of 8 in methanol-d<sub>4</sub>
- Figure S37. HSQC spectrum of 8 in methanol-d4
- Figure S38. HMBC spectrum of 8 in methanol-d4
- Figure S39. ROESY spectrum of 8 in methanol-d4

Figure S40. HRESIMS spectrum of 8

Figure S41. Purification of CD8+T cells in vitro.

Figure S42. Compounds affected the anti-tumor activity of CD8<sup>+</sup> T cells in vitro.



Figure S2. <sup>13</sup>C NMR and DEPT spectrum of **1** in methanol-*d*<sub>4</sub>





Figure S6. HRESIMS spectrum of 1



















Figure S19. HRESIMS spectrum of 3



Figure S20. <sup>1</sup>H NMR spectrum of **4** in methanol-*d*<sub>4</sub>





🔏 🕱 Spectru	①本企 + · % · 注止 ← · (+, +, +) だ(前 Q, 回 回 回 前 Spectrum from YYN, wi42 (sample 55) - M4.55. Experiment 1, +DA TOF MS (100 - 2000) from 3.918 min															40 4		
	1.0e5								357.1680								&	
Intensity, cps	5.0e4								379.1508									
	0.0-0	235.1330								358.1732 395.1255				441.1225				
	0.0e0 -	200	220	240	260	280	300	320	340 Mass/Charge, Da	360	380	400	420	440	460	480		
														r an				
Found elemental compositions Find Any Find MS Details MSMS Details Compound Details														ш_				
Ht	Formula	m/z	RDB ppm MS Rank	MSMS MSMS	Found	e cluster details	Charge +1	<ul> <li>MS result</li> </ul>	It summary for C21H25C	5. [M+H]+								
1	C21H24O5	357.1697 1	10.0 -4.6 1		NA/NA 0	Use m/z	% intensity Width 100.0 0.009	0 100%	<sup>1</sup>	357,1680	)							
							>	805	· ·	7/1								
					Eleme	Elements trom 60				60% -								
					Mass	tolerance (ppm)	10	405	202					358.1732 0.4 ppm				
					Intens	Intensity tolerance (%) 100				20/								
						#Cläheteroatoms greater than 0 Ion type: [M+H]+ V V <u>5 additional ions</u>												

Figure S25. HRESIMS spectrum of 4



Figure S26. CD spectrum of 4 in MeOH



Figure S28. <sup>13</sup>C NMR and DEPT spectrum of 7 in methanol-d<sub>4</sub>









Figure S33. HRESIMS spectrum of 7

















Figure S41. Purification of CD8+T cells in vitro.



Figure S42. Compounds affected the anti-tumor activity of CD8<sup>+</sup> T cells in vitro. Cells were incubated in the corresponding concentration of compounds or DMSO for 18 h and the fluorescence intensity was detected by a microplate reader (emission light 476nm, excitation light 514nm).