

SUPPORTING INFORMATION

Tetrabutylammonium Iodide-Catalyzed Oxidative α -Azidation of β -Ketocarbonyl Compounds Using Sodium Azide

Christopher Mairhofer¹, David Naderer¹, and Mario Waser^{1*}

*Email: mario.waser@jku.at

¹Institute of Organic Chemistry, Johannes Kepler University Linz, Altenbergerstr. 69, 4040 Linz, Austria

Table of Contents

1.	General Information	2
2.	Handling of Azido Compounds.....	3
3.	Synthesis of Substrates	4
4.	Oxidative α -Azidation Reactions	5
	General Procedure	5
	Characterization of the α -Azidation Products	5
5.	Oxidative α -Nitration Reactions.....	16
	General Procedure	16
	Characterization of the α -Nitration Products	16
6.	NMR Spectra	18
7.	Infrared Spectra	43
8.	High Resolution Mass Spectra	55

1. General Information

Nuclear magnetic resonance (NMR) spectra were recorded on a Bruker Avance III 300 MHz spectrometer with a broad band observe probe and a sample changer for 16 samples, on a Bruker Avance DRX 500 MHz spectrometer, and on a Bruker Avance III 700 MHz spectrometer with an Ascend magnet and TCI cryoprobe, which are all property of the Austro Czech NMR Research Center “RERI uasb”. All NMR spectra were referenced on the solvent residual peak (CDCl_3 : $\delta = 7.26 \text{ ppm}$ for ^1H -NMR, $\delta = 77.16 \text{ ppm}$ for ^{13}C -NMR, ^{19}F -NMR stayed unreferenced). ^1H -NMR and ^{19}F -NMR spectra are reported as follows: chemical shift (δ / ppm) (multiplicity, coupling constants, number of protons). Peak multiplicities are denoted as: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad, dd = doublet of doublet, etc. Infrared (IR) spectra were recorded on a Bruker Alpha II FTIR spectrometer with diamond ATR-module using OPUS software package and are reported in $\bar{\nu} / \text{cm}^{-1}$. High resolution mass spectra (HRMS) were recorded on an Agilent QTOF 6520 with an ESI source, owned and operated by the Institute of Analytical and General Chemistry, JKU. Melting points (MP) are recorded using a Büchi M-560 apparatus and are reported uncorrected in $\vartheta_m / ^\circ\text{C}$. Thin layer chromatography (TLC) was performed on Macherey-Nagel pre-coated TLC plates (silica gel, 60 F₂₅₄, 0.20 mm, ALUGRAM® Xtra SIL). TLC plates were visualized by irradiation with UV light at $\lambda_{\max} = 254 \text{ nm}$ or using permanent staining with permanganate. Preparative column chromatography was carried out using Davisil LC 60A 70–200 MICRON silica gel. The term *in vacuo* refers to removal of solvents by rotary evaporation (40°C , 950–20 mbar) followed by drying at high vacuum. All chemicals were purchased from commercial suppliers and used without further purification unless otherwise stated.

Anhydrous dibenzoyl peroxide (DBPO) was prepared from the commercially available 70–75% hydrate:¹ To a solution of wet dibenzoyl peroxide (10 g, 75%, remainder water, Sigma-Aldrich) in 100 mL of CH_2Cl_2 was added anhydrous Na_2SO_4 (10 g) at 0°C . The suspension was stirred for 30 min and then filtered through a sintered glass funnel. The solvent was removed *in vacuo* (water bath temperature $<40^\circ\text{C}$) and dried under high vacuum to obtain a free-flowing white solid which was stored at 4°C under argon.


Deactivated silica gel was prepared from the commercially available Davisil LC 60A 70–200 MICRON silica gel: To a suspension of silica gel (300 g) in 400 mL of Et_2O , triethylamine (10.0 mL) was added. Then, the solvent was removed *in vacuo* and dried under high vacuum to obtain a free-flowing white solid which was stored on the bench.

2. Handling of Azido Compounds²

Sodium azide (NaN_3 , CAS: 26628-22-8) is acutely toxic ($LD_{50, \text{mouse}} = 27 \text{ mg kg}^{-1}$) and should be handled only under adequate safety precautions. Sodium azide reacts with acids to form hydrazoic acid (HN_3) which is both toxic ($LD_{50, \text{mouse}} = 22 \text{ mg kg}^{-1}$) and a spontaneously explosive gas. Excess NaN_3 adhered on any surface (flasks, paper, etc.) should be destroyed in a fume hood by soaking with acidified *sodium nitrite* or by oxidation with *cerium(IV) ammonium nitrate*.^{2g} Organic azides are potentially explosive chemicals (PECs) that decompose with introduction of external energy (heat, light, shock and pressure). Azides with a $(\text{C} + \text{O}) / \text{N}$ ratio of < 1 are potentially explosive and should never be isolated. Azides with a $(\text{C} + \text{O}) / \text{N}$ ratio between 1 and 3 can be isolated but should be stored as solutions ($\leq 1 \text{ M}$) below room temperature with less than 5 grams of material. Azides with a $(\text{C} + \text{O}) / \text{N}$ ratio of ≥ 3 are normally stable and can be isolated and stored in pure form. Any azide synthesized should be stored below room temperature and in the dark.


(2) (a) Bräse, S.; Banert, K. *Organic Azides: Syntheses and Applications*; Wiley, 2010. DOI: 10.1002/9780470682517. (b) Bräse, S.; Gil, C.; Knepper, K.; Zimmermann, V. Organic azides: an exploding diversity of a unique class of compounds. *Angew. Chem. Int. Ed* **2005**, *44* (33), 5188–5240. (c) Treitler, D. S.; Leung, S. How Dangerous Is Too Dangerous? A Perspective on Azide Chemistry. *J. Org. Chem.* **2022**, *87* (17), 11293–11295. (d) Kolb, H. C.; Finn, M. G.; Sharpless, K. B. Click Chemistry: Diverse Chemical Function from a Few Good Reactions. *Angew. Chem. Int. Ed.* **2001**, *40* (11), 2004–2021. (e) Wiss, J.; Fleury, C.; Heuberger, C.; Onken, U.; Glor, M. Explosion and Decomposition Characteristics of Hydrazoic Acid in the Gas Phase. *Org. Process Res. Dev.* **2007**, *11*, 1096–1103. (f) Kopach, M. E.; Murray, M. M.; Braden, T. M.; Kobierski, M. E.; Williams, O. L. Improved Synthesis of 1-(Azidomethyl)-3,5-bis-(trifluoromethyl)benzene: Development of Batch and Microflow Azide Processes. *Org. Process Res. Dev.* **2009**, *13*, 152–160. (g) Bretherick, L. *Handbook of Reactive Chemical Hazards*, 4th ed.; Butterworths: London, 1990; p 1360.

3. Synthesis of Substrates



Indanone based ketoesters:

- X = OtBu, R = H: **1a**
- X = OMe, R = H: **1b**
- X = Oallyl, R = H: **1c**
- X = OBn, R = H: **1d**
- X = OAd, R = H: **1e**
- X = OCum, R = H: **1f**
- X = OtBu, R = 4-F: **1g**
- X = OtBu, R = 4-Me: **1h**
- X = OtBu, R = 5-F: **1i**
- X = OtBu, R = 5-Cl: **1j**
- X = OtBu, R = 5-Br: **1k**
- X = OtBu, R = 5-OMe: **1l**
- X = OtBu, R = 6-OMe: **1m**
- X = OtBu, R = 6-Me: **1n**



Lactone based ketoesters:

- X = OMe: **11a**
- X = Oallyl: **11b**
- X = Oallyl: **11c**

Indanone based diketones:

- X = Me, R = H: **1p**
- X = Ph, R = H: **1q**

Indanone based ketoamides:

- X = Nmorph, R = H: **1r**

All substrates are known compounds and prepared by following the literature procedures:

The indanone based *tert*-butyl β -ketoesters **1a**^{3,4}, **1g-n**^{3,4} were prepared starting from 1-indanone, CAS 83-33-0 or the respective commercially available 1-indanone derivatives using Boc-pyrrole, CAS 5176-27-2 as transfer-reagent. The other indanone based β -ketoesters **1b**⁵, **1c**⁶, were prepared starting from 1-indanone and their respective commercially available dialkyl carbonates. The indanone based β -ketoesters **1d-f**⁷ as well as the β -ketoamide **1r**⁸ were prepared starting from methyl 1-indanone-2-carboxylate **1b**. The indanone based 1,3-diketones **1p-q**^{3,4} were prepared according to a very similar procedure, but using imidazole based transfer reagents. These 1-acyl-imidazoles^{4,9} which are prepared from imidazole, CAS 288-32-4 and the respective commercially available acyl chlorides for diketones or chloroformates for β -ketoesters. Since these 1-acyl-imidazole reagents are more reactive than Boc-pyrrole, the reaction can be carried out at room temperature and is also a powerful alternative to prepare the β -ketoesters **1a-c**. The lactone based β -ketoesters **11a-c**^{10,11,12} were prepared from γ -butyrolactone, CAS 96-48-0 and the respective commercially available dialkyl carbonates or acyl chlorides, whereas also in this case the use of the previously described 1-acyl-imidazole transfer reagents turned out to be beneficial in terms of yield and selectivity.

[3] Moss, T. A.; Fenwick, D. R.; Dixon, D. J. Enantio- and Diastereoselective Catalytic Alkylation Reactions with Aziridines. *J. Am. Chem. Soc.* **2008**, *130*, 10076–10077.

[4] Y.-N. Duan, L.-Q. Cui, L.-H. Zuo, C. Zhang, Recyclable Hypervalent-Iodine-Mediated Dehydrogenative a,b'-Bifunctionalization of b-Keto Esters Under Metal-Free Conditions. *Chem. Eur. J.* **2015**, *21*, 13052–13057.

[5] Vita, M. V.; Waser, J. Azidation of β -Keto Esters and Silyl Enol Ethers with a Benziodoxole Reagent. *Org. Lett.* **2013**, *15* (13), 3246–3249.

[6] Craig, R. A.; Loskot, S. A.; Mohr, J. T.; Behenna, D. C.; Harned, A. M.; Stoltz, B. M. Palladium-Catalyzed Enantioselective Decarboxylative Allylic Alkylation of Cyclopentanones. *Org. Lett.* **2015**, *17* (21), 5160–5163.

[7] (a) Christoffers, J.; Önal, N., Azeotropic Transesterification of β -Keto Esters. *Eur. J. Org. Chem.* **2000**, 1633–1635. (b) Baumhof, P.; Mazitschek, R.; Giannis, A., A Mild and Effective Method for the Transesterification of Carboxylic Acid Esters. *Angew. Chem. Int. Ed.* **2001**, *40*, 3672–3674. (c) Lian, M.; Li, Z.; Du, J.; Meng, Q.; Gao, Z. Asymmetric Direct α -Hydroxylation of β -Oxo Esters by Phase-Transfer Catalysis Using Chiral Quaternary Ammonium Salts. *Eur. J. Org. Chem.* **2010**, *34*, 6525–6530. (d) Pericas, À.; Shafir, A.; Vallribera, A. Zinc(II) oxide: An efficient catalyst for selective transesterification of β -ketoesters. *Tetrahedron* **2008**, *64* (39), 9258–9263.

[8] Li, M.; Hawkins, A.; Barber, D. M.; Bultinck, P.; Herrebout, W.; Dixon, D. J. Enantio- and diastereoselective palladium catalysed arylative and vinylative allene carbocyclisation cascades. *Chem. Commun.* **2013**, *49* (46), 5265–5267.

[9] Q. Tian, Z. Gan, X. Wang, D. Li, W. Luo, H. Wang, Z. Dai, J. Yuan; Imidazolium Chloride: An Efficient Catalyst for Transamidation of Primary Amines, *Molecules* **2018**, 2234–2237.


[10] L. Cicco, V. Addante, A. Temperini, C. A. Donau, K. Karaghiosoff, F. M. Perna, V. Capriati, Toward Customized Tetrahydropyran Derivatives through Regioselective α -Lithiation and Functionalization of 2-Phenyltetrahydropyran, *Eur. J. Org. Chem.* **2016**, *19*, 3157.

[11] H. Kawada, H. Ebiike, M. Tsukazaki, M. Nakamura, K. Morikami, K. Yoshinari, M. Yoshida, K. Ogawa, N. Shimma, T. Tsukuda et al., Lead optimization of a dihydropyrrrolopyrimidine inhibitor against phosphoinositide 3-kinase (PI3K) to improve the phenol glucuronic acid conjugation *Bioorg. Med. Chem. Lett.* **2013**, *23*, 673.

[12] T. Ooi, T. Miki, K. Fukumoto, K. Maruoka, Asymmetric Synthesis of α -Acyl- γ -butyrolactones Possessing All-Carbon Quaternary Stereocenters by Phase-Transfer-Catalyzed Alkylation, *Adv. Synth. Catal.* **2006**, *348*, 1539.

4. Oxidative α -Azidation Reactions


General Procedure



Sodium azide (7.9 mg, 120 μ mol, 1.2 eq) and TBAI (7.5 mg, 20 μ mol, 20 mol%) were suspended in a stirred solution (900-1000 rpm) of the respective substrate (100 μ mol, 1.00 eq) in 1.0 mL of DCE at r.t. Then, a solution of anhydrous benzoyl peroxide (29.1 mg, 120 μ mol, 1.2 eq) in 1.0 mL of DCE was added to the suspension and stirred for 20 h.

The reaction solution was then diluted with 8 mL dichloromethane and extracted with 5 mL of sat. aq. NaHCO₃. The aqueous phase was then extracted twice with 10 mL of DCM. The organic layer and the extracts were then filtered consecutively through a pad of anhydr. sodium sulfate and deactivated silica gel. The solvents were removed *in vacuo*. In most cases the products were already obtained in high purity (> 95 %) after this work up. If necessary further purification can be achieved by silica gel column chromatography.

Characterization of the α -Azidation Products



tert-Butyl 2-azido-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2a):^{13,14,15,16,17} Obtained in 94% yield (25.7 mg, 94.0 μ mol). *cf.*: 1.0 mmol scale, 94% yield (256.9 mg, 940.0 μ mol). This compound was purified by column chromatography on silica gel (eluent: heptanes/EtOAc = 19:1). Yellowish-white solid; **¹H-NMR** (300 MHz, CDCl₃, 298 K, δ / ppm): 7.82 (d, J = 7.7 Hz, 1H), 7.66 (t, J = 7.5 Hz, 1H), 7.48 - 7.39 (m, 2H), 3.64 (d, J = 17.2 Hz, 1H), 2.99 (d, J = 17.2 Hz, 1H), 1.45 (s, 9H); **¹³C-NMR** (75 MHz, CDCl₃, 298 K, δ / ppm): 198.1, 167.4, 152.3, 136.4, 133.3, 128.4, 126.5, 125.6, 84.6, 70.6, 38.6, 28.0; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹): 2984, 2928, 2853, 2110, 1747, 1736, 1718, 1604, 1589, 1548, 1466, 1431, 1397, 1372, 1353, 1326, 1271, 1259, 1215, 1145, 1091, 1054, 1027, 961, 913, 871, 844, 834, 818, 804, 756, 729, 711, 688, 661, 623, 598, 561, 533, 459, 416; **HRMS** (ESI⁺-QqTOF, *m/z*): calculated for C₁₄H₁₆N₃O₃ [M+H]⁺: 274.1186, found: 274.1186;

[13] M. Tiffner, L. Stockhammer, J. Schögenhumer, K. Röser and M. Waser; Towards an Asymmetric Organocatalytic α -Azidation of β -Ketoesters *Molecules* **2018**, 1142-1151.


[14] M. Uyanik, N. Sahara, M. Tsukahara, Y. Hattori and K. Ishihara; Chemo- and Enantioselective Oxidative α -Azidation of Carbonyl Compounds; *Angew. Chem. Int. Ed.* **2020**, 59, 17110–17117.

[15] Q.-H. Deng, T. Bleith, H. Wadeohl, and L. H. Gade Enantioselective Iron-Catalyzed Azidation of β -Keto Esters and Oxindoles *J. Am. Chem. Soc.* **2013**, 135, 5356–5359


[16] M. V. Vita and J. Waser; Azidation of β -Keto Esters and Silyl Enol Ethers with a Benziodoxole Reagent; *Org. Lett.* **2013**, 15, 3246-3249

[17] K. Shibatomi, Y. Soga, A. Narayama, I. Fujisawa, and S. Iwasa; Highly Enantioselective Chlorination of β -Ketoesters and Subsequent S_N2 Displacement of Tertiary Chlorides: A Flexible Method for the Construction of Quaternary Stereogenic Centers; *J. Am. Chem. Soc.* **2012**, 134, 9836–9839


calculated for $C_{14}H_{19}N_4O_3 [M+NH_4]^+$: 291.1452, found: 291.1452 (major); calculated for $C_{14}H_{15}N_3O_3Na [M+Na]^+$: 296.1006, found: 296.1006; calculated for $C_{14}H_{15}N_3O_3K [M+K]^+$: 312.0745, found: 312.0745, **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.64; **MP** (uncorrected, ϑ_m / °C): 65.0 - 67.5.




Methyl 2-azido-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2b):^{14,16} Obtained in 96% yield (22.2 mg, 96.0 μmol); white solid; **1H-NMR** (500 MHz, CDCl₃, 298 K, δ / ppm): 7.83 (d, J = 7.7 Hz, 1H), 7.69 (t, J = 7.5 Hz, 1H), 7.55 - 7.41 (m, 2H), 3.81 (s, 3H), 3.68 (d, J = 17.4 Hz, 1H), 3.04 (d, J = 17.3 Hz, 1H); **13C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 197.5, 169.1, 152.2, 136.6, 133.1, 128.6, 126.6, 125.8, 70.3, 53.7, 38.6; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹): 3475, 3409, 3187, 3082, 3040, 3010, 2956, 2922, 2851, 2449, 2428, 2326, 2278, 2178, 2135, 2099, 2081, 1744, 1710, 1601, 1587, 1479, 1467, 1440, 1427, 1344, 1326, 1305, 1278, 1218, 1186, 1161, 1122, 1098, 1050, 998, 965, 951, 903, 826, 806, 777, 739, 696, 626, 599, 550, 469, 437, 406; **HRMS** (ESI⁺-QqTOF, m/z): calculated for $C_{11}H_{10}N_3O_3 [M+H]^+$: 232.0717, found: 232.0717; calculated for $C_{11}H_{13}N_4O_3 [M+NH_4]^+$: 249.0982, found: 249.0982 (major), calculated for $C_{11}H_9N_3O_3Na [M+Na]^+$: 254.0536, found: 254.0539; calculated for $C_{11}H_9N_3O_3K [M+K]^+$: 270.0275, found: 270.0275; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.50; **MP** (uncorrected, ϑ_m / °C): 54.7 - 56.5.




Allyl 2-azido-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2c): Obtained in 98% yield (25.2 mg, 98.0 μmol); yellow solid; **1H-NMR** (500 MHz, CDCl₃, 298 K, δ / ppm): 7.84 (d, J = 7.7 Hz, 1H), 7.69 (t, J = 7.5 Hz, 1H), 7.47 (d, J = 7.8 Hz, 1H), 7.46 (t, J = 7.5 Hz, 1H), 5.85 (ddt, J = 17.2, 10.5, 5.6 Hz, 1H), 5.29 - 5.19 (m, 2H), 4.70 (tt, J = 5.8, 1.4 Hz, 2H), 3.69 (d, J = 17.3 Hz, 1H), 3.05 (d, J = 17.3 Hz, 1H); **13C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 197.4, 168.3, 152.2, 136.6, 133.2, 130.9, 128.6, 126.6, 125.8, 119.3, 70.4, 67.2, 38.7; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹): 3077, 2954, 2923, 2852, 2109, 1744, 1713, 1649, 1605, 1588, 1464, 1426, 1377, 1362, 1328, 1265, 1227, 1213, 1179, 1124, 1089, 1046, 994, 938, 908, 865, 826, 806, 749, 697, 617, 599, 550, 468, 423; **HRMS** (ESI⁺-QqTOF, m/z): calculated for $C_{13}H_{12}N_3O_3 [M+H]^+$: 258.0873, found: 258.0874; calculated for $C_{13}H_{15}N_4O_3Na [M+NH_4]^+$: 275.1139, found: 275.1139 (major), calculated for $C_{13}H_{11}N_3O_3K [M+K]^+$: 296.0432, found: 296.0432; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.55; **MP** (uncorrected, ϑ_m / °C): 80.1 - 84.7.




Benzyl 2-azido-1-oxo-2,3-dihydro-1*H*-indene-2-carboxylate (2d): Obtained in 87% yield (26.7 mg, 87.0 μmol); yellow oil; **$^1\text{H-NMR}$** (500 MHz, CDCl_3 , 298 K, δ / ppm): 7.84 (d, $J = 7.9$ Hz, 1H), 7.68 (t, $J = 7.4$ Hz, 1H), 7.49 - 7.43 (m, 2H), 7.36 - 7.31 (m, 3H), 7.28 - 7.24 (m, 2H), 5.30 (d, $J = 12.3$ Hz, 1H), 5.20 (d, $J = 12.3$ Hz, 1H), 3.66 (d, $J = 17.3$ Hz, 1H), 3.04 (d, $J = 17.3$ Hz, 1H); **$^{13}\text{C-NMR}$** (126 MHz, CDCl_3 , 298 K, δ / ppm): 197.4, 168.5, 152.1, 136.6, 134.8, 133.2, 128.8, 128.7, 128.6, 128.2, 126.6, 125.8, 70.4, 68.4, 38.6; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 3474, 3407, 3343, 3194, 3096, 3066, 3032, 2981, 2930, 2848, 2681, 2638, 2607, 2485, 2442, 2359, 2317, 2110, 1954, 1920, 1742, 1710, 1607, 1583, 1491, 1465, 1437, 1392, 1370, 1339, 1292, 1260, 1243, 1212, 1168, 1151, 1132, 1099, 1050, 1026, 961, 924, 904, 892, 866, 839, 798, 749, 730, 700, 653, 617, 601, 561, 548, 497, 468, 441, 416; **HRMS** (ESI $^+$ -QqTOF, m/z): calculated for $\text{C}_{17}\text{H}_{14}\text{N}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 308.1030, found: 308.1032; calculated for $\text{C}_{17}\text{H}_{17}\text{N}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 325.1295, found: 325.1295 (major); calculated for $\text{C}_{17}\text{H}_{13}\text{N}_3\text{O}_3\text{Na}$ [$\text{M}+\text{Na}]^+$: 330.0849, found: 330.0849; calculated for $\text{C}_{17}\text{H}_{13}\text{N}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 346.0588, found: 346.0588; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.55.




Adamantyl 2-azido-1-oxo-2,3-dihydro-1*H*-indene-2-carboxylate (2e):^{13,16} Obtained in 77% yield (27.1 mg, 77.0 μmol); colourless oil; **$^1\text{H-NMR}$** (500 MHz, CDCl_3 , 298 K, δ / ppm): 7.81 (d, $J = 7.7$ Hz, 1H), 7.66 (t, $J = 7.6$ Hz, 1H), 7.55 - 7.38 (m, 2H), 3.64 (d, $J = 17.1$ Hz, 1H), 2.98 (d, $J = 17.1$ Hz, 1H), 2.15 (s, 3H), 2.07 (d, $J = 3.0$ Hz, 6H), 1.63 (t, $J = 2.8$ Hz, 6H); **$^{13}\text{C-NMR}$** (126 MHz, CDCl_3 , 298 K, δ / ppm): 198.2, 167.1, 152.4, 136.3, 133.4, 128.4, 126.4, 125.6, 84.7, 70.6, 41.2, 38.7, 36.1, 31.0; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 2911, 2853, 2110, 1742, 1714, 1607, 1589, 1457, 1426, 1355, 1264, 1230, 1213, 1182, 1155, 1103, 1090, 1046, 963, 912, 862, 835, 806, 749, 722, 698, 636, 549, 469, 419; **HRMS** (ESI $^+$ -QqTOF, m/z): calculated for $\text{C}_{20}\text{H}_{22}\text{N}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 352.1656, found: 352.1653; calculated for $\text{C}_{20}\text{H}_{26}\text{N}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 369.1921, found: 369.1922 (major); calculated for $\text{C}_{20}\text{H}_{22}\text{N}_3\text{O}_3\text{Na}$ [$\text{M}+\text{Na}]^+$: 374.1475, found: 374.1478; calculated for $\text{C}_{20}\text{H}_{22}\text{N}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 390.1214, found: 390.1213; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.65.




2-Phenylpropan-2-yl 2-azido-1-oxo-2,3-dihydro-1*H*-indene-2-carboxylate (2f):¹³ Obtained in 61% yield (20.5 mg, 61.0 µmol); yellow oil; **¹H-NMR** (500 MHz, CDCl₃, 298 K, δ / ppm): 7.85 (d, *J* = 7.7 Hz, 1H), 7.68 (t, *J* = 7.6 Hz, 1H), 7.52 - 7.43 (m, 1H), 7.33 - 7.27 (m, 1H), 7.27 - 7.20 (m, 2H), 3.67 (d, *J* = 17.2 Hz, 1H), 3.03 (d, *J* = 17.2 Hz, 1H), 1.81 (s, 3H), 1.77 (s, 3H); **¹³C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 197.9, 166.7, 152.2, 144.5, 136.4, 133.4, 128.5, 128.5, 127.6, 126.5, 125.6, 124.3, 85.5, 70.6, 38.6, 28.7, 28.1; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹): 2983, 2933, 2111, 1746, 1715, 1606, 1588, 1496, 1465, 1449, 1428, 1385, 1368, 1264, 1231, 1214, 1133, 1101, 1044, 1030, 939, 909, 835, 809, 751, 697, 628, 592, 551, 468, 420; **HRMS** (ESI⁺-QqTOF, *m/z*): calculated for C₁₉H₂₁N₄O₃ [M+NH₄]⁺: 353.1608, found: 353.1608 (major); calculated for C₁₉H₁₇N₃O₃Na [M+Na]⁺: 358.1162, found: 358.1163; calculated for C₁₉H₁₇N₃O₃K [M+K]⁺: 374.0901, found: 374.0901; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, *R_f* / 1): 0.57.




tert-Butyl 2-azido-4-fluoro-1-oxo-2,3-dihydro-1*H*-indene-2-carboxylate (2g): Obtained in 89% yield (25.9 mg, 89.0 µmol); yellow oil; **¹H-NMR** (500 MHz, CDCl₃, 298 K, δ / ppm): 7.63 (d, *J* = 7.5 Hz, 1H), 7.45 (td, *J* = 7.8, 4.5 Hz, 1H), 7.35 (t, *J* = 8.3 Hz, 1H), 3.64 (d, *J* = 17.5 Hz, 1H), 2.98 (d, *J* = 17.5 Hz, 1H), 1.46 (s, 9H); **¹³C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 197.1 (d, *J* = 2.8 Hz), 167.0, 159.7 (d, *J* = 251.6 Hz), 138.2 (d, *J* = 19.5 Hz), 136.0 (d, *J* = 5.0 Hz), 130.4 (d, *J* = 6.4 Hz), 122.5 (d, *J* = 20.0 Hz), 121.3 (d, *J* = 4.0 Hz), 85.0, 70.3, 34.6, 28.0; **¹⁹F-NMR** (471 MHz, CDCl₃, 298 K, δ / ppm): -118.29 (dd, *J* = 8.6, 4.5 Hz, 1F); **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹): 2981, 2933, 2116, 1746, 1722, 1619, 1594, 1482, 1458, 1426, 1395, 1371, 1339, 1264, 1244, 1186, 1147, 1048, 1035, 992, 868, 833, 813, 774, 746, 717, 686, 580, 551, 500, 469, 463; **HRMS** (ESI⁺-QqTOF, *m/z*): calculated for C₁₄H₁₈FN₄O₃ [M+NH₄]⁺: 309.1357, found: 309.1357 (major); calculated for C₁₄H₁₄FN₃O₃Na [M+Na]⁺: 314.0911, found: 314.0911; calculated for C₁₄H₁₄FN₃O₃K [M+K]⁺: 330.0651, found: 330.0651; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, *R_f* / 1): 0.67.




tert-Butyl 2-azido-4-methyl-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2h): Obtained in 91% yield (26.1 mg, 91.0 μmol); yellowish residue; **$^1\text{H-NMR}$** (500 MHz, CDCl_3 , 298 K, δ / ppm): 7.65 (d, $J = 7.6$ Hz, 1H), 7.47 (d, $J = 7.3$ Hz, 1H), 7.35 (t, $J = 7.5$ Hz, 1H), 3.52 (d, $J = 17.2$ Hz, 1H), 2.86 (d, $J = 17.2$ Hz, 1H), 2.33 (s, 3H), 1.47 (s, 9H); **$^{13}\text{C-NMR}$** (126 MHz, CDCl_3 , 298 K, δ / ppm): 198.3, 167.6, 151.3, 136.9, 135.8, 133.1, 128.6, 123.0, 84.6, 70.5, 37.6, 28.0, 17.9; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 3079, 2980, 2932, 2870, 2444, 2361, 2297, 2112, 1742, 1715, 1606, 1592, 1479, 1458, 1421, 1395, 1370, 1336, 1264, 1250, 1201, 1148, 1050, 1027, 953, 895, 859, 839, 830, 813, 769, 746, 721, 691, 661, 626, 574, 550, 509, 461; **HRMS** ($\text{ESI}^+ \text{-QqTOF}$, m/z): calculated for $\text{C}_{15}\text{H}_{18}\text{N}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 288.1343, found: 288.1343; calculated for $\text{C}_{15}\text{H}_{21}\text{N}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 305.1608, found: 305.1608 (major); calculated for $\text{C}_{15}\text{H}_{17}\text{N}_3\text{O}_3\text{Na}$ [$\text{M}+\text{Na}]^+$: 310.1162, found: 310.1162; calculated for $\text{C}_{15}\text{H}_{17}\text{N}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 326.0901, found: 326.0901; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.67; **MP** (uncorrected, ϑ_m / $^\circ\text{C}$): 65.0 - 67.5.




tert-Butyl 2-azido-5-fluoro-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2i):¹³ Obtained in 93% yield (27.1 mg, 93.0 μmol); yellowish solid; **$^1\text{H-NMR}$** (500 MHz, CDCl_3 , 298 K, δ / ppm): 7.82 (dd, $J = 8.3, 5.2$ Hz, 1H), 7.19 - 7.04 (m, 2H), 3.62 (d, $J = 17.4$ Hz, 1H), 2.96 (d, $J = 17.4$ Hz, 1H), 1.45 (s, 9H); **$^{13}\text{C-NMR}$** (126 MHz, CDCl_3 , 298 K, δ / ppm): 196.2, 168.1 (d, $J = 259.3$ Hz), 167.1, 155.4 (d, $J = 10.5$ Hz), 129.7 (d, $J = 1.8$ Hz), 128.0 (d, $J = 10.7$ Hz), 116.9 (d, $J = 23.8$ Hz), 113.3 (d, $J = 22.8$ Hz), 84.8, 70.7, 38.5 (d, $J = 2.2$ Hz), 28.0; **$^{19}\text{F-NMR}$** (471 MHz, CDCl_3 , 298 K, δ / ppm): -99.4 (dt, $J = 8.5, 5.1$ Hz, 1F); **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 3432, 3102, 3073, 2988, 2925, 2853, 2675, 2426, 2289, 2242, 2107, 1920, 1796, 1734, 1720, 1614, 1593, 1483, 1455, 1432, 1398, 1371, 1335, 1298, 1253, 1192, 1148, 1086, 1046, 942, 906, 900, 867, 830, 802, 787, 759, 738, 691, 655, 629, 620, 551, 503, 476, 438; **HRMS** ($\text{ESI}^+ \text{-QqTOF}$, m/z): calculated for $\text{C}_{14}\text{H}_{15}\text{FN}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 309.1357, found: 309.1357; calculated for $\text{C}_{14}\text{H}_{18}\text{FN}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 309.1357, found: 309.1357 (major); calculated for $\text{C}_{14}\text{H}_{14}\text{FN}_3\text{O}_3\text{Na}$ [$\text{M}+\text{Na}]^+$: 314.0911, found: 314.0911; calculated for $\text{C}_{14}\text{H}_{14}\text{FN}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 330.0651, found: 330.0651; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.67; **MP** (uncorrected, ϑ_m / $^\circ\text{C}$): 62.0 - 65.0




tert-Butyl 2-azido-5-chloro-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2j): Obtained in 78% yield (24.0 mg, 78.0 μmol); yellow solid; **$^1\text{H-NMR}$** (500 MHz, CDCl_3 , 298 K, δ / ppm): 7.74 (d, $J = 8.2$ Hz, 1H), 7.46 (s, 1H), 7.42 (d, $J = 8.2$ Hz, 1H), 3.61 (d, $J = 17.4$ Hz, 1H), 2.96 (d, $J = 17.4$ Hz, 1H), 1.46 (s, 9H); **$^{13}\text{C-NMR}$** (126 MHz, CDCl_3 , 298 K, δ / ppm): 196.8, 167.1, 153.7, 143.1, 131.8, 129.3, 126.7, 126.6, 84.9, 70.6, 38.3, 28.0; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 3093, 3065, 2989, 2928, 2854, 2359, 2326, 2293, 2164, 2116, 1918, 1742, 1716, 1599, 1578, 1456, 1423, 1396, 1370, 1323, 1297, 1279, 1261, 1245, 1210, 1146, 1069, 1048, 914, 868, 834, 797, 787, 755, 735, 722, 688, 612, 550, 524, 480, 451, 435; **HRMS** ($\text{ESI}^+ \text{-QqTOF}$, m/z): calculated for $\text{C}_{14}\text{H}_{15}^{35}\text{ClN}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 308.0796, found: 308.0799; calculated for $\text{C}_{14}\text{H}_{18}^{35}\text{ClN}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 325.1062, found: 325.1061 (major); calculated for $\text{C}_{14}\text{H}_{18}^{37}\text{ClN}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 327.1037, found: 327.1038; calculated for $\text{C}_{14}\text{H}_{14}^{35}\text{ClN}_3\text{O}_3\text{Na}$ [$\text{M}+\text{Na}]^+$: 330.0616, found: 330.0616; calculated for $\text{C}_{14}\text{H}_{14}^{35}\text{ClN}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 346.0355, found: 346.0355; calculated for $\text{C}_{14}\text{H}_{14}^{37}\text{ClN}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 348.0331, found: 348.0331; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.67; **MP** (uncorrected, ϑ_m / $^\circ\text{C}$): 76.0 - 78.9.




tert-Butyl 2-azido-5-bromo-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2k):^{13,15} Obtained in 96% yield (33.8 mg, 96.0 μmol); white solid; **$^1\text{H-NMR}$** (500 MHz, CDCl_3 , 298 K, δ / ppm): 7.67 (d, $J = 8.2$ Hz, 1H), 7.64 (s, 1H), 7.58 (d, $J = 8.2$ Hz, 1H), 3.61 (d, $J = 17.4$ Hz, 1H), 2.96 (d, $J = 17.4$ Hz, 1H), 1.46 (s, 9H); **$^{13}\text{C-NMR}$** (126 MHz, CDCl_3 , 298 K, δ / ppm): 197.0, 167.0, 153.8, 132.2, 132.1, 132.0, 129.8, 126.7, 84.9, 70.5, 38.2, 28.0; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 3089, 3002, 2981, 2926, 2852, 2493, 2308, 2162, 2111, 1816, 1736, 1718, 1592, 1468, 1455, 1418, 1396, 1371, 1322, 1299, 1280, 1244, 1212, 1196, 1177, 1143, 1212, 1196, 1177, 1143, 1131, 1101, 1058, 1046, 1036, 913, 868, 847, 834, 792, 749, 730, 713, 687, 615, 596, 550, 517, 477, 460, 439, 424, 406; **HRMS** ($\text{ESI}^+ \text{-QqTOF}$, m/z): calculated for $\text{C}_{14}\text{H}_{15}^{79}\text{BrN}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 352.0291, found: 352.0291; calculated for $\text{C}_{14}\text{H}_{15}^{81}\text{BrN}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 354.0272, found: 354.0272; calculated for $\text{C}_{14}\text{H}_{18}^{79}\text{BrN}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 369.0557, found: 369.0557; calculated for $\text{C}_{14}\text{H}_{18}^{81}\text{BrN}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 371.0538, found: 369.0538 (major); calculated for $\text{C}_{14}\text{H}_{14}^{79}\text{BrN}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 389.9850, found: 389.9850; calculated for $\text{C}_{14}\text{H}_{14}^{81}\text{BrN}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 391.9831, found: 391.9831; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.68; **MP** (uncorrected, ϑ_m / $^\circ\text{C}$): 91.3 - 93.6.




tert-Butyl 2-azido-5-methoxy-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2l):¹⁵ Obtained in 90% yield (27.3 mg, 90.0 µmol); yellow oil; **1H-NMR** (500 MHz, CDCl₃, 298 K, δ / ppm): 7.73 (d, J = 8.5 Hz, 1H), 6.94 (d, J = 8.9 Hz, 1H), 6.86 (s, 1H), 3.89 (s, 3H), 3.57 (d, J = 17.2 Hz, 1H), 2.91 (d, J = 17.1 Hz, 1H), 1.45 (s, 9H); **13C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 195.9, 167.7, 166.6, 155.5, 127.3, 126.3, 116.5, 109.6, 84.4, 70.9, 55.9, 38.6, 28.0; **IR** (neat, FT-ATR, 298 K, ν / cm⁻¹): 3476, 3406, 3191, 3096, 3066, 3031, 2981, 2931, 2845, 2680, 2633, 2606, 2485, 2441, 2361, 2316, 2185, 2111, 2059, 1920, 1742, 1710, 1607, 1582, 1491, 1465, 1437, 1392, 1370, 1339, 1293, 1259, 1244, 1208, 1144, 1132, 1098, 1052, 1025, 925, 904, 891, 866, 839, 797, 756, 729, 703, 653, 617, 601, 561, 547, 478, 462, 441; **HRMS** (ESI⁺-QqTOF, m/z): calculated for C₁₅H₁₈N₃O₄ [M+H]⁺: 304.1292, found: 304.1292; calculated for C₁₅H₂₁N₄O₄ [M+NH₄]⁺: 321.1557, found: 321.1557 (major); calculated for C₁₅H₁₇N₃O₄Na [M+Na]⁺: 326.1111, found: 326.1111; calculated for C₁₅H₁₇N₃O₄K [M+K]⁺: 342.0851, found: 342.0852; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.50.




tert-Butyl 2-azido-6-methyl-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2m):¹³ Obtained in 96% yield (27.6 mg, 96.0 µmol); yellow oil; **1H-NMR** (500 MHz, CDCl₃, 298 K, δ / ppm): 7.59 (s, 1H), 7.47 (d, J = 7.8 Hz, 1H), 7.33 (d, J = 7.8 Hz, 1H), 3.57 (d, J = 17.1 Hz, 1H), 2.93 (d, J = 17.1 Hz, 1H), 2.41 (s, 3H), 1.44 (s, 9H); **13C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 198.1, 167.5, 149.8, 138.5, 137.6, 133.4, 126.1, 125.4, 84.5, 70.9, 38.3, 27.9, 21.2; **IR** (neat, FT-ATR, 298 K, ν / cm⁻¹): 2980, 2932, 2870, 2358, 2325, 2286, 2110, 1742, 1715, 1617, 1584, 1494, 1457, 1424, 1395, 1276, 1241, 1202, 1147, 1102, 1047, 1102, 1047, 1102, 1046, 1036, 950, 905, 889, 839, 825, 798, 789, 748, 731, 695, 626, 584, 551, 503, 476, 453, 439, 417; **HRMS** (ESI⁺-QqTOF, m/z): calculated for C₁₅H₁₈N₃O₃ [M+H]⁺: 288.1343, found: 288.1343; calculated for C₁₅H₂₁N₄O₃ [M+NH₄]⁺: 305.1608, found: 305.1608 (major); calculated for C₁₅H₁₇N₃O₃Na [M+Na]⁺: 310.1162, found: 310.1163; calculated for C₁₅H₁₇N₃O₃K [M+K]⁺: 326.0901, found: 326.0907; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.67.




tert-Butyl 2-azido-6-methoxy-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (2n):^{13,15} Obtained in 78% yield (23.7 mg, 78.0 µmol); yellow oil; **1H-NMR** (500 MHz, CDCl₃, 298 K, δ / ppm): 7.33 (d, J = 8.4 Hz, 1H), 7.24 (dd, J = 8.4, 2.6 Hz, 1H), 7.21 (d, J = 2.6 Hz, 1H), 3.84 (s, 3H), 3.54 (d, J = 16.9 Hz, 1H), 2.90 (d, J = 16.8 Hz, 1H), 1.44 (s, 9H); **13C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 198.0, 167.4, 160.0, 145.3, 134.5, 127.1, 125.8, 106.4, 84.5, 71.3, 55.8, 38.0, 27.9; **IR** (neat, FT-ATR, 298 K, ν / cm⁻¹): 3403, 3299, 3067, 2993, 2972, 2852, 2727, 2680, 2622, 2576, 2445, 2422, 2359, 2272, 2110, 2022, 1912, 1735, 1711, 1613, 1585, 1547, 1492, 1460, 1432, 1395, 1370, 1341, 1304, 1277, 1234, 1211, 1190, 1157, 1128, 1092, 1056, 1033, 1022, 959, 895, 876, 841, 824, 801, 787, 772, 739, 690, 634, 589, 555, 539, 524, 480, 463, 450, 435, 423; **HRMS** (ESI⁺-QqTOF, m/z): calculated for C₁₅H₁₈N₃O₄ [M+H]⁺: 304.1292, found: 304.1291; calculated for C₁₅H₂₁N₄O₄ [M+NH₄]⁺: 321.1557, found: 321.1557 (major); calculated for C₁₅H₁₇N₃O₄Na [M+Na]⁺: 326.1111, found: 326.1111; calculated for C₁₅H₁₇N₃O₄K [M+K]⁺: 342.0851, found: 342.0851; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.60.




2-Acetyl-2-azido-2,3-dihydro-1H-inden-1-one (5a): Obtained in 61% yield (13.1 mg, 61.0 µmol); yellow oil; **1H-NMR** (700 MHz, CDCl₃, 298 K, δ / ppm): 7.82 (d, J = 7.7 Hz, 1H), 7.70 (t, J = 7.5 Hz, 1H), 7.52 (d, J = 7.9 Hz, 1H), 7.47 (t, J = 7.5 Hz, 1H), 3.71 (d, J = 17.4 Hz, 1H), 3.08 (d, J = 17.4 Hz, 1H), 2.29 (s, 3H); **13C-NMR** (176 MHz, CDCl₃, 298 K, δ / ppm): 200.8, 197.8, 152.2, 136.8, 133.5, 128.8, 126.7, 125.7, 77.8, 37.1, 26.1; **IR** (neat, FT-ATR, 298 K, ν / cm⁻¹): 3412, 3077, 2954, 2926, 2854, 2362, 2342, 2324, 2100, 1726, 1708, 1606, 1588, 1465, 1424, 1356, 1334, 1301, 1256, 1211, 1188, 1154, 1090, 1053, 1025, 982, 959, 901, 877, 813, 778, 739, 704, 671, 644, 583, 564, 549, 524, 502, 468, 426; **HRMS** (ESI-QqTOF, m/z): calculated for C₁₁H₉N₃O₂ [M+H]⁺: 216.0768, found: 216.0770; calculated for C₁₁H₁₃N₄O₂ [M+NH₄]⁺: 233.1033, found: 233.1033 (major); calculated for C₁₂H₁₀N₃O₄ [M+HCOO]⁺: 260.0677, found: 260.0684; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.43.




2-Azido-2-benzoyl-2,3-dihydro-1H-inden-1-one (5b): Obtained in 83% yield (23.0 mg, 83.0 μmol); yellow oil; **1H-NMR** (700 MHz, CDCl_3 , 298 K, δ / ppm): 7.92 (d, $J = 7.7$ Hz, 1H), 7.85 (d, $J = 8.2$ Hz, 2H), 7.72 (d, $J = 7.4$ Hz, 1H), 7.57 (t, $J = 7.4$ Hz, 1H), 7.52 (t, $J = 7.6$ Hz, 1H), 7.49 (d, $J = 7.6$ Hz, 1H), 7.41 (t, $J = 7.8$ Hz, 2H), 3.77 (d, $J = 17.5$ Hz, 1H), 3.25 (d, $J = 17.5$ Hz, 1H); **13C-NMR** (176 MHz, CDCl_3 , 298 K, δ / ppm): 197.9, 194.5, 151.2, 136.6, 134.0, 133.9, 133.5, 129.2, 128.9, 128.9, 126.9, 126.0, 76.3, 39.2; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 3068, 2925, 2853, 2449, 2280, 2099, 1786, 1763, 1716, 1681, 1598, 1465, 1448, 1429, 1303, 1257, 1234, 1213, 1155, 1089, 1032, 994, 961, 946, 899, 871, 846, 799, 777, 741, 701, 688, 671, 610, 550, 468, 443, 419; **HRMS** (ESI-QqTOF, m/z): calculated for $\text{C}_{16}\text{H}_{12}\text{N}_3\text{O}_2$ [$\text{M}+\text{H}]^+$: 278.0924, found: 278.0924; calculated for $\text{C}_{16}\text{H}_{15}\text{N}_4\text{O}_2$ [$\text{M}+\text{NH}_4]^+$: 295.1190, found: 295.1190 (major); calculated for $\text{C}_{16}\text{H}_{11}\text{N}_3\text{O}_2\text{Na}$ [$\text{M}+\text{Na}]^+$: 300.0743, found: 300.0743; calculated for $\text{C}_{16}\text{H}_{11}\text{N}_3\text{O}_2\text{K}$ [$\text{M}+\text{K}]^+$: 316.0483, found: 316.0483; calculated for $\text{C}_{17}\text{H}_{12}\text{N}_3\text{O}_4$ [$\text{M}+\text{HCOO}]^-$: 322.0833, found: 322.0833; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.48.




2-Azido-2-(morpholine-4-carbonyl)-2,3-dihydro-1H-inden-1-one (6): Obtained in 78% yield (22.3 mg, 78.0 μmol); yellow solid; **1H-NMR** (700 MHz, CDCl_3 , 298 K, δ / ppm): 7.84 (d, $J = 7.4$ Hz, 1H), 7.67 (t, $J = 7.5$ Hz, 1H), 7.48 - 7.43 (m, 2H), 3.81 - 3.54 (m, 9H), 3.17 (d, $J = 16.9$ Hz, 1H); **13C-NMR** (176 MHz, CDCl_3 , 298 K, δ / ppm): 197.5, 165.8, 150.3, 136.5, 133.6, 129.2, 128.8, 126.6, 125.7, 72.4, 66.9, 38.6; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 2964, 2922, 2856, 2360, 2249, 2099, 1718, 1638, 1607, 1558, 1457, 1425, 1361, 1330, 1301, 1270, 1255, 1233, 1213, 1184, 1155, 1113, 1090, 1065, 1010, 942, 914, 873, 852, 836, 808, 795, 759, 732, 703, 674, 646, 626, 582, 550, 534, 478, 454, 418; **HRMS** (ESI⁺-QqTOF, m/z): calculated for $\text{C}_{14}\text{H}_{15}\text{N}_4\text{O}_3$ [$\text{M}+\text{H}]^+$: 287.1139, found: 287.1139 (major); calculated for $\text{C}_{14}\text{H}_{14}\text{N}_4\text{O}_3\text{Na}$ [$\text{M}+\text{Na}]^+$: 309.0958, found: 309.0958; calculated for $\text{C}_{14}\text{H}_{14}\text{N}_4\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 325.0697, found: 325.0694; **TLC** (silica gel K60, 200 μm , F254, heptanes/IPA = 7:3, 298 K, R_f / 1): 0.36; **MP** (uncorrected, ϑ_m / $^\circ\text{C}$): 125.0 - 127.0.



Methyl 3-azido-2-oxotetrahydrofuran-3-carboxylate (7a): Obtained in 55% yield (10.2 mg, 55.0 µmol); yellowish oil; **1H-NMR** (700 MHz, CDCl₃, 298 K, δ / ppm): 4.45 (2 × ddd, J = 8.9, 7.7, 6.1 Hz, 2H), 3.91 (s, 3H), 2.77 (ddd, J = 13.3, 7.6, 5.5 Hz, 1H), 2.29 (ddd, J = 13.6, 7.9, 6.6 Hz, 1H); **13C-NMR** (176 MHz, CDCl₃, 298 K, δ / ppm): 170.7, 167.4, 66.6, 66.5, 54.2, 33.6; **IR** (neat, FT-ATR, 298 K, ν / cm⁻¹): 2960, 2925, 2852, 2360, 2114, 1775, 1748, 1669, 1600, 1559, 1483, 1451, 1437, 1380, 1318, 1245, 1216, 1171, 1118, 1019, 996, 955, 918, 841, 803, 791, 745, 705, 687, 634, 615, 601, 552, 536, 469, 455, 434; **HRMS** (ESI⁺-QqTOF, m/z): calculated for C₆H₈N₃O₄ [M+H]⁺: 186.0509, found: 186.0508; calculated for C₆H₈N₃O₄ [M+NH₄]⁺: 203.0775, found: 203.0776 (major); **TLC** (silica gel K60, 200 µm, F254, heptanes/IPA = 7:3, 298 K, R_f / 1): 0.34.




Allyl 3-azido-2-oxotetrahydrofuran-3-carboxylate (7b): Obtained in 74% yield (15.6 mg, 74.0 µmol); yellowish oil; **1H-NMR** (700 MHz, CDCl₃, 298 K, δ / ppm): 5.93 (ddt, J = 16.6, 10.3, 5.8 Hz, 1H), 5.39 (dd, J = 17.1, 1.4 Hz, 1H), 5.33 (dd, J = 10.4, 1.3 Hz, 1H), 4.85 - 4.74 (m, 2H), 4.53 - 4.40 (m, 2H), 2.78 (ddd, J = 13.3, 7.6, 5.4 Hz, 1H), 2.30 (ddd, J = 13.5, 8.0, 6.6 Hz, 1H); **13C-NMR** (176 MHz, CDCl₃, 298 K, δ / ppm): 170.7, 166.6, 130.5, 120.3, 67.9, 66.6, 66.5, 33.6; **IR** (neat, FT-ATR, 298 K, ν / cm⁻¹): 3090, 2990, 2956, 2928, 2862, 2115, 1775, 1747, 1650, 1483, 1447, 1424, 1379, 1316, 1270, 1241, 1211, 1171, 1117, 1020, 975, 937, 843, 790, 740, 714, 688, 661, 640, 597, 551, 535; **HRMS** (ESI⁺-QqTOF, m/z): calculated for C₈H₉N₃O₄ [M+H]⁺: 212.0666, found: 212.0666; calculated for C₈H₁₃N₄O₄ [M+NH₄]⁺: 229.0931, found: 229.0931 (major); calculated for C₈H₉N₃O₄Na [M+Na]⁺: 234.0485, found: 234.0488; calculated for C₈H₉N₃O₄K [M+K]⁺: 250.0225, found: 250.0223; **TLC** (silica gel K60, 200 µm, F254, heptanes/IPA = 7:3, 298 K, R_f / 1): 0.57.



3-Azido-3-benzoyldihydrofuran-2(3H)-one (7c):¹⁸ Obtained in 69% yield (16.0 mg, 69.0 μmol); yellowish oil; **$^1\text{H-NMR}$** (700 MHz, CDCl_3 , 298 K, δ / ppm): 5.93 (ddt, $J = 16.6, 10.3, 5.8$ Hz, 1H), 5.39 (dd, $J = 17.1, 1.4$ Hz, 1H), 5.33 (dd, $J = 10.4, 1.3$ Hz, 1H), 4.85 - 4.74 (m, 2H), 4.53 - 4.40 (m, 2H), 2.78 (ddd, $J = 13.3, 7.6, 5.4$ Hz, 1H), 2.30 (ddd, $J = 13.5, 8.0, 6.6$ Hz, 1H); **$^{13}\text{C-NMR}$** (176 MHz, CDCl_3 , 298 K, δ / ppm): 170.7, 166.6, 130.5, 120.3, 67.9, 66.6, 66.5, 33.6; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm^{-1}): 3067, 2983, 2923, 2858, 2355, 2107, 1771, 1673, 1596, 1579, 1482, 1449, 1377, 1308, 1272, 1243, 1214, 1176, 1102, 1021, 1001, 953, 886, 844, 796, 760, 718, 686, 650, 616, 579, 540, 499, 418; **HRMS** (ESI⁺-QqTOF, m/z): calculated for $\text{C}_{11}\text{H}_{10}\text{N}_3\text{O}_3$ [$\text{M}+\text{H}]^+$: 232.0717, found: 232.0717; calculated for $\text{C}_{11}\text{H}_{13}\text{N}_4\text{O}_3$ [$\text{M}+\text{NH}_4]^+$: 249.0982, found: 287.0982 (major); calculated for $\text{C}_{11}\text{H}_9\text{N}_3\text{O}_3\text{Na}$ [$\text{M}+\text{Na}]^+$: 254.0536, found: 254.0536; calculated for $\text{C}_{11}\text{H}_9\text{N}_3\text{O}_3\text{K}$ [$\text{M}+\text{K}]^+$: 270.0275, found: 270.0276; **TLC** (silica gel K60, 200 μm , F254, heptanes/IPA = 7:3, 298 K, R_f / 1): 0.51.

5. Oxidative α -Nitration Reactions


General Procedure




Sodium nitrite (15.2 mg, 220 μmol , 2.2 eq) and TBAI (7.4 mg, 20 μmol , 20 mol%) were suspended in a stirred solution (900-1000 rpm) of the respective substrate (100 μmol , 1.00 eq) in 1.0 mL of DCE at r.t. Then, a solution of anhydrous benzoyl peroxide (29.1 mg, 120 μmol , 1.2 eq) in 1.0 mL of DCE was added to the suspension and stirred for 20 h.

The reaction solution was then diluted with 8 mL dichloromethane and extracted with 5 mL of sat. aq. NaHCO₃. The aqueous phase was then extracted twice with 10 mL of DCM. The organic layer and the extracts were then filtered consecutively through a pad of anhydr. sodium sulfate and deactivated silica gel. The solvents were removed *in vacuo*. In most cases the products were already obtained in high purity (> 95 %) after this work up. If necessary further purification can be achieved by silica gel column chromatography.


Characterization of the α -Nitration Products



tert-Butyl 2-nitro-1-oxo-2,3-dihydro-1H-indene-2-carboxylate (10a): Obtained in 84% yield (23.3 mg, 94.0 μmol); white solid; **¹H-NMR** (700 MHz, CDCl₃, 298 K, δ / ppm): 7.86 (d, J = 7.7 Hz, 1H), 7.71 (t, J = 7.5 Hz, 1H), 7.53 (d, J = 7.7 Hz, 1H), 7.48 (t, J = 7.5 Hz, 1H), 4.11 (d, J = 17.9 Hz, 1H), 3.99 (d, J = 17.9 Hz, 1H), 1.49 (s, 9H); **¹³C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 188.4, 162.0, 150.1, 137.0, 132.9, 129.1, 126.5, 126.2, 96.7, 86.1, 37.5, 27.8.; **IR** (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹): 2984, 2930, 2878, 2854, 1748, 1719, 1656, 1604, 1589, 1548, 1465, 1431, 1396, 1371, 1353, 1325, 1272, 1260, 1215, 1145, 1091, 1056, 1026, 961, 912, 871, 844, 834, 818, 803, 755, 730, 711, 688, 661, 625, 598, 561, 533, 459, 414; **HRMS** (ESI⁺-QqTOF, *m/z*): calculated for C₁₄H₁₆NO₅ [M+H]⁺: 278.1023, found: 278.1024; calculated for C₁₄H₁₉N₂O₅ [M+NH₄]⁺: 295.1288, found: 295.1288 (major); calculated for C₁₄H₁₅NO₅Na [M+Na]⁺: 300.0842, found: 300.0842; calculated for C₁₄H₁₅NO₅K [M+K]⁺: 316.0582, found: 316.0582; **TLC** (silica gel K60, 200 μm , F254, heptanes/EtOAc = 7:3, 298 K, R_f / 1): 0.47; **MP** (uncorrected, ϑ_m / °C): 75.9 – 78.4.



Methyl 2-nitro-1-oxo-2,3-dihydro-1*H*-indene-2-carboxylate (10b)¹⁹: Obtained in 86% yield (19.9 mg, 96.0 µmol); colourless oil; **¹H-NMR** (700 MHz, CDCl₃, 298 K, δ / ppm): 7.87 (d, *J* = 7.8 Hz, 1H), 7.73 (t, *J* = 7.5 Hz, 1H), 7.54 (d, *J* = 7.8 Hz, 1H), 7.50 (t, *J* = 7.6 Hz, 1H), 4.15 (d, *J* = 18.0 Hz, 1H), 4.05 (d, *J* = 18.0 Hz, 1H), 3.89 (s, 3H); **¹³C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 188.0, 163.8, 150.0, 137.3, 132.7, 129.3, 126.5, 126.4, 96.1, 54.5, 37.5; **IR** (neat, FT-ATR, 298 K, *ν* / cm⁻¹): 2959, 1755, 1729, 1605, 1552, 1434, 1264, 1215, 1202, 1181, 1091, 1023, 949, 903, 894, 798, 761, 689, 663, 632, 467; **HRMS** (ESI⁺-QqTOF, *m/z*): calculated for C₁₁H₁₀NO₅ [M+H]⁺: 236.0553, found: 236.0553; calculated for C₁₁H₁₃N₂O₅ [M+NH₄]⁺: 253.0820, found: 253.0820 (major); calculated for C₁₁H₉NO₅Na [M+Na]⁺: 258.0373, found: 258.0373; calculated for C₁₁H₉NO₅K [M+K]⁺: 274.0112, found: 274.0112; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, *R_f* / 1): 0.29.




Allyl 2-nitro-1-oxo-2,3-dihydro-1*H*-indene-2-carboxylate (10c): Obtained in 75% yield (19.3 mg, 75.0 µmol); yellowish oil; **¹H-NMR** (700 MHz, CDCl₃, 298 K, δ / ppm): 7.88 (d, *J* = 7.7 Hz, 1H), 7.73 (t, *J* = 7.5 Hz, 1H), 7.54 (d, *J* = 7.7 Hz, 1H), 7.50 (t, *J* = 7.5 Hz, 1H), 5.89 (ddt, *J* = 16.5, 11.0, 5.7 Hz, 1H), 5.38 - 5.27 (m, 2H), 4.81 - 4.74 (m, 2H), 4.16 (d, *J* = 18.0 Hz, 1H), 4.05 (d, *J* = 18.0 Hz, 1H); **¹³C-NMR** (126 MHz, CDCl₃, 298 K, δ / ppm): 188.0, 163.0, 150.0, 137.2, 132.7, 130.3, 129.3, 126.5, 126.5, 120.1, 96.1, 68.2, 37.5; **IR** (neat, FT-ATR, 298 K, *ν* / cm⁻¹): 3063, 2940, 2453, 2324, 1920, 1782, 1757, 1731, 1656, 1599, 1557, 1466, 1450, 1423, 1350, 1296, 1275, 1255, 1217, 1178, 1134, 1093, 1070, 1034, 1019, 995, 941, 926, 902, 840, 809, 794, 755, 692, 616, 601, 555, 500, 467, 437, 416; **HRMS** (ESI⁺-QqTOF, *m/z*): calculated for C₁₃H₁₂N₁O₅ [M+H]⁺: 262.0710, found: 262.0710; calculated for C₁₃H₁₅N₂O₅ [M+NH₄]⁺: 279.0975, found: 279.0975 (major); calculated for C₁₃H₁₁NO₅Na [M+Na]⁺: 284.0529, found: 284.0529; calculated for C₁₃H₁₁NO₅K [M+K]⁺: 300.0269, found: 300.0272; **TLC** (silica gel K60, 200 µm, F254, heptanes/EtOAc = 7:3, 298 K, *R_f* / 1): 0.62.


[19] F.-Z. Han, L.-L. Li, L.-N. Jia, X.-P. Hu; Catalyst-free nitration of the aliphatic CAH bonds of tertiary *b*-keto esters with tert-butyl nitrite: Access to *a*-quaternary *a*-amino acid precursors; *Tetrahedron Lett.* 2022, 99, 153844-153849

6. NMR Spectra


2a, ^1H -NMR (300 MHz, CDCl_3 , 298 K, δ / ppm):




2a, ^{13}C -NMR (126 MHz, CDCl_3 , 298 K, δ / ppm):




2b, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2b, ^{13}C -NMR (126 MHz, CDCl_3 , 298 K, δ / ppm):




2c, ¹H-NMR (500 MHz, CDCl₃, 298 K, δ / ppm):




2c, ^{13}C -NMR (126 MHz, CDCl_3 , 298 K, δ / ppm):




2d, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2d, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2e, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2e, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2f, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2f, ^{13}C -NMR (126 MHz, CDCl_3 , 298 K, δ / ppm):




2g, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2g, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2g, $^{19}\text{F-NMR}$ (471 MHz, CDCl_3 , 298 K, δ / ppm):




2h, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2h, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2i, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2i, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2i, $^{19}\text{F-NMR}$ (471 MHz, CDCl_3 , 298 K, δ / ppm):




2j, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2j, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2k, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2k, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2l, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2l, ^{13}C -NMR (126 MHz, CDCl_3 , 298 K, δ / ppm):




2m, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2m, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




2n, $^1\text{H-NMR}$ (500 MHz, CDCl_3 , 298 K, δ / ppm):




2n, $^{13}\text{C-NMR}$ (126 MHz, CDCl_3 , 298 K, δ / ppm):




5a, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




5a, $^{13}\text{C-NMR}$ (176 MHz, CDCl_3 , 298 K, δ / ppm):




5b, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




5b, ¹³C-NMR (176 MHz, CDCl₃, 298 K, δ / ppm):




6, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




6, $^{13}\text{C-NMR}$ (176 MHz, CDCl_3 , 298 K, δ / ppm):




7a, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




7a, ^{13}C -NMR (176 MHz, CDCl_3 , 298 K, δ / ppm):




7b, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




7b, ¹³C-NMR (176 MHz, CDCl₃, 298 K, δ / ppm):




7c, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




7c, $^{13}\text{C-NMR}$ (176 MHz, CDCl_3 , 298 K, δ / ppm):




10a, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




10a, $^{13}\text{C-NMR}$ (176 MHz, CDCl_3 , 298 K, δ / ppm):




10b, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




10b, $^{13}\text{C-NMR}$ (176 MHz, CDCl_3 , 298 K, δ / ppm):



10c, $^1\text{H-NMR}$ (700 MHz, CDCl_3 , 298 K, δ / ppm):




10c, ^{13}C -NMR (176 MHz, CDCl_3 , 298 K, δ / ppm):




7. Infrared Spectra


2a, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):




2b, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):




2c, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2d, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2e, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):




2f, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):




2g, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2h, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2i, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2j, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2k, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2l, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




2m, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):




2n, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):




5a, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




5b, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




6, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




7a, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




7b, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




7c, IR (neat, FT-ATR, 298 K, $\bar{\nu}$ / cm⁻¹):




10a, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):



10b, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):



10c, IR (neat, FT-ATR, 298 K, \bar{v} / cm⁻¹):




8. High Resolution Mass Spectra

(major only)


2a, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₄H₁₉N₄O₃ [M+NH₄]⁺: 291.1452, found: 291.1452 (0.00 ppm)




2b, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₁H₁₃N₄O₃ [M+NH₄]⁺: 249.0982, found: 249.0982 (0.00 ppm)




2c, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₃H₁₅N₄O₃Na [M+NH₄]⁺: 275.1139, found: 275.1139 (0.00 ppm)




2d, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₇H₁₇N₄O₃ [M+NH₄]⁺: 325.1295, found: 325.1295 (0.00 ppm)




2e, IR HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₂₀H₂₆N₄O₃ [M+NH₄]⁺: 369.1921, found: 369.1922 (0.27 ppm)




2f, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₉H₂₁N₄O₃ [M+NH₄]⁺: 353.1608, found: 353.1608 (0.00 ppm)




2g, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₄H₁₈FN₄O₃ [M+NH₄]⁺: 309.1357, found: 309.1357 (0.00 ppm)




2h, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₅H₂₁N₄O₃ [M+NH₄]⁺: 305.1608, found: 305.1608 (0.00 ppm)




2i, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₄H₁₈FN₄O₃ [M+NH₄]⁺: 309.1357, found: 309.1357 (0.00 ppm)




2j, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₄H₁₈³⁵ClN₄O₃ [M+NH₄]⁺: 325.1062, found: 325.1061 (-0.31 ppm)




2k, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₄H₁₈⁸¹BrN₄O₃ [M+NH₄]⁺: 371.0538, found: 369.0538 (0.00 ppm)




2l, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₅H₂₁N₄O₄ [M+NH₄]⁺: 321.1557, found: 321.1557 (0.00 ppm)




2m, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₅H₂₁N₄O₃ [M+NH₄]⁺: 305.1608, found: 305.1608 (0.00 ppm)




2n, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₅H₂₁N₄O₄ [M+NH₄]⁺: 321.1557, found: 321.1557 (0.00 ppm)




5a, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₁H₁₃N₄O₂ [M+NH₄]⁺: 233.1033, found: 233.1033 (0.00 ppm)




5b, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₆H₁₅N₄O₂ [M+NH₄]⁺: 295.1190, found: 295.1190 (0.00 ppm)




6, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₄H₁₅N₄O₃ [M+H]⁺: 287.1139, found: 287.1139 (0.00 ppm)




7a, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₆H₈N₃O₄ [M+NH₄]⁺: 203.0775, found: 203.0776 (0.49 ppm)




7b, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₈H₁₃N₄O₄ [M+NH₄]⁺: 229.0931, found: 229.0931 (0.00 ppm)




7c, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₁H₁₃N₄O₃ [M+NH₄]⁺: 249.0982, found: 287.0982 (0.00 ppm)




10a, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₄H₁₉N₂O₅ [M+NH₄]⁺: 295.1288, found: 295.1288 (0.00 ppm)



10b, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₁H₁₃N₂O₅ [M+NH₄]⁺: 253.0820, found: 253.0820 (0.00 ppm)



10c, HRMS (ESI⁺-QqTOF, *m/z*):

calculated for C₁₃H₁₅N₂O₅ [M+NH₄]⁺: 279.0975, found: 279.0975 (0.00 ppm)

